

Supplementary Data

Supplemental Table 1. Search strategy for MEDLINE

<p>“follow-up” OR cohort OR survival OR mortality OR "all-cause mortality"</p>	<p>AND</p>	<p>dairy OR milk OR yogurt OR cheese OR kefir OR butter OR "dairy products"</p>	<p>AND</p>	<p>meta OR review</p>
--	------------	---	------------	-------------------------------

Supplementary Data**Supplemental Table 2.** Covariates considered in the analysis of the cohorts included in the meta-analysis.

Author	Covariates included in the analyses	Meta-analyses including each paper
Kahn et al. (S1)	Age, sex, smoking, history of vascular disease, hypertension, and diabetes	12, 13, 15, 18
Mann et al. (S2)	Age, sex, smoking, and social class	12-19
Ness et al. (S3)	Age, smoking, BP, cholesterol, BMI, forced expiratory volume, social class, education, deprivation, siblings, car user, angina, ECG ischemia, bronchitis, and alcohol	12-15, 17, 18
Elwood et al. (S4)	Age, smoking, social class, IHD, BMI, energy, alcohol, fasting cholesterol, HDL cholesterol and triglycerides	12-15, 17, 18
Trichopoulou et al. (S5)	Age, gender, smoking, education, weight and height, hip circum., insulin, hypertension, hypercholesterolemia, and food groups	12, 17
van der Pols et al. (S6)	Age, sex, area, energy, fruit, vegetables, eggs, protein, fat, and energy	12
Fortes et al. (S7)	Age, sex, education, BMI, smoking, cognitive function, and chronic diseases	13, 16-19
Knoops et al. (S8)	Age, sex, alcohol, physical activity, smoking, number of years of education, BMI, chronic diseases, and study center	13, 17, 18
Paganini-Hill et al. (S9)	Age, sex, smoking, exercise, BMI, alcohol, hypertension, angina, MI, stroke, diabetes, rheumatoid arthritis, and cancer	13-15, 17, 18
Engberink et al. (S10)	Age, sex, BMI, SBP, total cholesterol, family history of MI, use of estrogens, smoking, educational level, alcohol consumption, total energy, saturated fat, and intake of fruit, vegetables, meat, fish, coffee, and tea	13, 18
Bonthuis et al. (S11)	Age, sex, total energy intake, body mass index, alcohol intake, school leaving age, physical activity level, pack years of smoking, dietary supplement use, b-carotene treatment during trial, presence of any medical condition, and dietary calcium intake	13-17, 19
Goldbohm et al. (S12)	Age; education; cigarette, cigar, or pipe smoking; non-occupational physical activity; occupational physical activity; BMI; multivitamin use; alcohol; energy; energy-adjusted mono- and polyunsaturated fat intakes; and vegetable and fruit consumption	13-16, 19

Supplementary Data

Author, country	Covariates included in the analyses	Meta-analyses including each paper
Dalmeijer et al. (S13)	Gender, age, total energy intake, physical activity, smoking, education, BMI, ethanol, coffee, fruit, vegetables, fish, meat and bread	13
Soedamah-Muthu et al. (S14)	Age, ethnicity, employment grade, smoking, alcohol intake, BMI, physical activity, family history of CHD/ hypertension, and fruit, vegetable, bread, meat, fish, coffee, tea and total energy intake	13-15, 17, 19
van Aerde et al. (S15)	Age, sex, BMI, smoking, educational level, total energy intake, alcohol consumption, physical activity and intake of meat, fish, bread, vegetables, fruit, coffee, and tea	13, 15, 17, 19
Huang et al. (S16)	Age, gender, BMI, region, ethnicity, education level, marriage, history of disease (cardiovascular disease and/or cancer), smoking, drinking, chewing betel nut, supplement use, overall Dietary Index–Revised (dairy score excluded), calcium intake, and vitamin D intake	13, 17
Michaelsson et al. (S17)	Age, body mass index, height, total energy intake, total alcohol intake, healthy dietary pattern, calcium and vitamin D supplementation, historic use of cortisone, educational level, living alone, physical activity level estimated as metabolic equivalents, smoking status, and Charlson’s comorbidity index; in women, use of estrogen replacement therapy and nulliparity	13-15, 17
Praagman et al. (S18)	Age, sex, total energy intake, smoking habit, BMI, physical activity, education level, hypertension at baseline, intakes of alcohol and energy, and adjusted intakes of fruit and vegetables	13, 19
Wang et al. (S19)	Age categories, smoking status, drinking status, physical activity, sleeping duration, body mass index, education level, participation in health check-ups, green-leafy vegetable intake, and history of hypertension, diabetes, and liver disease	13, 14, 17
Dik et al. (S20)	Age, sex, center, smoking, pre-diagnostic BMI, tumor sub-site (colon or rectum), disease stage, differentiation grade, and total energy intake	14
Yang et al. (S21)	Age, sex, tumor stage and folate and total energy intake	14

Supplementary Data

Author, country	Covariates included in the analyses	Meta-analyses including each paper
Fraser et al. (S22)	Age, sex, BMI, smoking, cognitive status, medical history, and education	16, 19
Whiteman et al. (S23)	Age, sex, smoking, alcohol, and SES	16
Sluik et al. (S24)	Age and center-stratified; adjusted for gender, education, underlying dietary patterns, alcohol consumption, smoking, physical activity and prevalence of heart disease, cancer or stroke.	17
Boggs et al. (S25)	Age, each DASH component, total energy intake, education, marital status, physical activity, TV watching, smoking, and alcohol	17
Bongard et al. (S26)	Age, center, income, obesity, alcohol, smoking, physical activity, chronic disease, and diet quality score	17
Fraser et al. (S27)	Age, smoking, and physical activity	17, 19
Hays et al. (S28)	Age, living alone, poverty, smoking, alcohol, cognitive status, and self-rated health	17
Kouris-Blazos et al. (S29)	Age, energy intake, sex, smoking, and ethnic origin	17
Kurotani et al. (S30)	Age; sex; public health center area; BMI; smoking; physical activity; history of hypertension, diabetes or dyslipidemia; coffee and green tea consumption; and occupation	17
Martinez-Gonzalez et al. (S31)	Age, years of university education, BMI, smoking, physical activity, TV watching, history of depression, and baseline hypertension, hypercholesterolemia, energy intake, egg consumption, potato consumption, and adoption of special diets, in addition to all the other items in the Mediterranean Diet Score	17
Prinelli et al. (S32)	Age, sex, education, BMI, physical activity, smoking, time spent TV watching, and energy intake	17
Trichopoulou et al. (S33)	Age, sex, education, smoking, waist-to-hip ratio, BMI, physical activity, and energy intake	17
Vormund, 2015, Switzerland (S34)	Age, sex, survey wave, marital status, smoking, BMI, region and nationality, mutually adjusted	17

Supplementary Data

Author, country	Covariates included in the analyses	Meta-analyses including each paper
Yu, 2015, USA (S35)	Age, race, enrollment source, income, marital status, medical insurance, smoking, physical activity, sitting time, energy intake, BMI, menopausal status, hormone therapy, disease status, education, and standardized HEI score without dairy	17
Yu, 2014, China (S36)	Age, education, income, smoking, alcohol, multivitamin, menopausal status and hormone therapy, physical activity, BMI, waist-to-hip ratio, history of cardiovascular disease, diabetes, or hypertension, and energy intake	17

BMI: body mass index; BP: blood pressure; CHD: coronary heart disease; DASH: disabilities of the arm, shoulder and hand; ECG: electrocardiography; HDL: high density lipoprotein; HEI: healthy eating index; IHD; ischemic heart disease; MI: myocardial infarction; SES; socioeconomic status; SBP: systolic blood pressure; TV: television.

Supplementary Data

Supplemental Table 3. Risk of bias assessed using the AMSTAR2 tool.

Author	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	Total	Risk of bias
Elwood et al. (12)	Y	Y	Y	Y	N	N	N	Y	N	N	Y	N	N	Y	N	Y	8	Acceptable
Guo et al. (13)	Y	Y	Y	Y	N	N	N	Y	Y	Y	Y	Y	Y	Y	Y	Y	13	Very good
Larsson et al. (14)	Y	Y	Y	Y	N	N	N	Y	N	N	Y	N	N	Y	N	Y	8	Acceptable
Mullie et al. (15)	Y	Y	Y	Y	N	N	N	Y	N	N	Y	N	N	Y	Y	Y	9	Good
O’Sullivan et al. (16)	Y	Y	Y	Y	Y	Y	N	Y	Y	N	Y	Y	Y	Y	Y	Y	14	Very good
Schwingshackl et al. (17)	Y	Y	Y	Y	Y	Y	N	Y	Y	N	Y	Y	Y	Y	Y	Y	14	Very good
Soedamah- Muthu et al. (18)	Y	Y	Y	Y	Y	Y	N	Y	N	N	Y	N	N	Y	Y	Y	12	Very good
Tong et al. (19)	Y	Y	Y	Y	N	N	N	Y	N	N	Y	N	N	Y	Y	Y	9	Good

Supplementary Data

Supplemental Table 4. Certainty of evidence in systematic reviews and meta-analyses for all-cause mortality outcome assessed using GRADE.

Reference	<i>n</i> , studies	Study design	Certainty assessment					<i>n</i> , participants		Effect		Certainty	Importance
			Risk of bias ¹	Inconsistency ²	Indirectness	Imprecision	Other considerations	<i>n</i> , total	<i>n</i> , cases (%)	RR (95% CI)	Absolute risk ⁹		
Elwood et al. (12)	6	Cohort studies	serious	very serious	serious ³	not serious	publication bias strongly suspected all plausible residual confounding would suggest spurious effect, while no effect was observed ⁷	24466	5092 (20.8%)	0.87 (0.77 to 0.98)	Less than 1 case per 1,000 population	⊕○○○ VERY LOW	CRITICAL
Guo et al. (13)	17	Cohort studies	not serious	serious	not serious	serious ⁵	all plausible residual confounding would suggest spurious effect, while no effect was observed dose response gradient	175063	21222 (12.1%)	0.99 (0.96 to 1.03)	Less than 1 case per 1,000 population	⊕⊕○○ LOW	CRITICAL
Larsson et al. (14)	13	Cohort studies	serious	serious	not serious	serious ⁶	publication bias strongly suspected all plausible residual confounding would suggest spurious effect, while no effect was observed dose response gradient ⁷	367505	81401 (22.1%)	not estimable		⊕○○○ VERY LOW	CRITICAL
Mullie et al. (15)	10	Cohort studies	serious	very serious	not serious	serious ⁵	publication bias strongly suspected all plausible residual confounding would suggest spurious effect, while no effect was observed dose response gradient ⁸	281788	63545 (22.6%)	1.01 (0.96 to 1.06)	Less than 1 case per 1,000 population	⊕○○○ VERY LOW	CRITICAL
O'Sullivan et al. (16)	6	Cohort studies	not serious	not serious	serious ⁴	serious ⁵	publication bias strongly suspected all plausible residual confounding would suggest spurious effect, while no effect was observed ⁸	43797	17272 (39.4%)	1.01 (0.92 to 1.11)	Less than 1 case per 1,000 population	⊕○○○ VERY LOW	CRITICAL

Supplementary Data


Reference	<i>n, studies</i>	Study design	Certainty assessment					<i>n, participants</i>		Effect		Certainty	Importance
			Risk of bias ¹	Inconsistency ²	Indirectness	Imprecision	Other considerations	<i>n, total</i>	<i>n, cases (%)</i>	RR (95% CI)	Absolute risk ⁹		
Schwingshackl et al. (17)	26	Cohort studies	not serious	very serious	serious ⁴	serious ⁵	publication bias strongly suspected all plausible residual confounding would suggest spurious effect, while no effect was observed dose response gradient ⁸	938817	126759 (13.5%)	1.03 (0.98 to 1.07)	Less than 1 case per 1,000 population	⊕○○○ VERY LOW	CRITICAL
Soedamah-Muthu et al. (18)	8	Cohort studies	not serious	serious	not serious	serious ⁵	all plausible residual confounding would suggest spurious effect, while no effect was observed dose response gradient	62779	23949 (38.1%)	0.99 (0.95 to 1.03)	Less than 1 case per 1,000 population	⊕⊕○○ LOW	CRITICAL
Tong et al. (19)	8	Cohort studies	serious	not serious	not serious	serious ⁵	all plausible residual confounding would suggest spurious effect, while no effect was observed dose response gradient	177657	21365 (12.0%)	1.03 (0.99 to 1.07)	Less than 1 case per 1,000 population	⊕⊕○○ LOW	CRITICAL

RR: risk ratio; CI: Confidence interval.

¹ Based on AMSTAR tool score. From 1 to 5: not serious; from 6 to 10: serious; and from 11 to 16: very serious; ² Based on I². From 0 to 40%: not serious; from 40 to 80%: serious; and from 80 to 100% and not reported: very serious; ³ The original studies are not aimed at measuring the association; ⁴ Dairy products consumption is not the main observation; ⁵ CI includes the null value; ⁶ No reported by the meta-analysis; ⁷ Does not assess publication bias or find publication bias by Egger test; ⁸ Find publication bias by Egger test; ⁹ For dairy products consumption.

Supplementary Data

Supplemental Figure 1. Forest plot for each dairy product consumption increment of 1 serving/day in adults.



RR: risk ratio; CI: confidence interval

Supplementary data

Supplemental References

- S1. Kahn HA, Phillips RL, Snowdon DA, Choi W. Association between reported diet and all-cause mortality. Twenty-one-year follow-up on 27,530 adult Seventh-Day Adventists. *Am J Epidemiol* 1984; 119:775–787.
- S2. Mann JI, Appleby PN, Key TJ, Thorogood M. Dietary determinants of ischaemic heart disease in health conscious individuals. *Heart* 1997; 78: 450–455.
- S3. Ness AR, Smith GD, Hart C. Milk, coronary heart disease and mortality. *J Epidemiol Community Health* 2001;55: 379–382.
- S4. Elwood PC, Pickering JE, Fehily AM, Hughes J, Ness AR. Milk drinking, ischaemic heart disease and ischaemic stroke I. Evidence from the Caerphilly cohort. *Eur J Clin Nutr* 2004; 58: 711–717.
- S5. Trichopoulou A, Psaltopoulou T, Orfanos P, Trichopoulos D. Diet and physical activity in relation to overall mortality amongst adult diabetics in a general population cohort. *J Intern Med* 2006; 259:583–591.
- S6. van der Pols JC, Gunnell D, Williams GM, Holly JM, Bain C, Martin RM. Childhood dairy and calcium intake and cardiovascular mortality in adulthood: 65-year follow-up of the Boyd Orr cohort. *Heart* 2009;95:1600–1606.
- S7. Fortes C, Forastiere F, Farchi S, Rapiti E, Pastori G, Perucci CA. Diet and overall survival in a cohort of very elderly people. *Epidemiology* 2000;11:440–5.

Supplementary Data

- S8. Knoops KT, Groot de LC, Fidanza F, Alberti-Fidanza A, Kromhout D, van Staveren WA. Comparison of three different dietary scores in relation to 10-year mortality in elderly European subjects: the HALE project. *Eur J Clin Nutr* 2006;60:746–55.
- S9. Paganini-Hill A, Kawas CH, Corrada MM. Non-alcoholic beverage and caffeine consumption and mortality: the Leisure World Cohort Study. *Prev Med* 2007;44:305–10.
- S10. Engberink MF, Soedaman-Muthu SS, Boessenkool-Pape J, van Rooij FJA, Hofman A, Witteman JCM, Geleijnse JM. Dairy intake in relation to all-cause mortality and risk of cardiovascular disease: the Rotterdam Study. American Heart Association, San Francisco. 2010.
- S11. Bonthuis M, Hughes MC, Ibiebele TI, Green AC, Van Der Pols JC. Dairy consumption and patterns of mortality of Australian adults. *Eur J Clin Nutr* 2010; 64: 569–577.
- S12. Goldbohm RA, Chorus AM, Galindo GF, Schouten LJ, van den Brandt PA. Dairy consumption and 10-y total and cardiovascular mortality: a prospective cohort study in the Netherlands. *Am J Clin Nutr* 2011; 93: 615–627.
- S13. Dalmeijer GW, Struijk EA, van der Schouw YT, Soedamah-Muthu SS, Verschuren W. M, Boer JM, Geleijnse JM, Beulens JW. Dairy intake and coronary heart disease or stroke – a population-based cohort study. *Int J Cardiol* 2013; 167: 925–929.
- S14. Soedamah-Muthu SS, Masset G, Verberne L, Geleijnse JM, Brunner EJ. Consumption of dairy products and associations with incident diabetes, CHD and mortality in the Whitehall II study. *Br J Nutr* 2013; 109: 718–726.

Supplementary Data

S15. van Aerde MA, Soedamah-Muthu SS, Geleijnse JM, Snijder MB, Nijpels G, Stehouwer CD, Dekker JM. Dairy intake in relation to cardiovascular disease mortality and all-cause mortality: the Hoorn Study. *Eur J Nutr* 2013; 52: 609–616.

S16. Huang LY, Wahlqvist ML, Huang YC, Lee MS. Optimal dairy intake is predicated on total, cardiovascular, and stroke mortalities in a Taiwanese cohort. *J Am Coll Nutr*. 2014;33:426–36.

S17. Michaelsson K, Wolk A, Langenskiöld S, Basu S, Warensjö Lemming E, Melhus H, Byberg L. Milk intake and risk of mortality and fractures in women and men: cohort studies. *BMJ*. 2014;349:g6015.

S18. Praagman J, Franco OH, Ikram MA, Soedamah-Muthu SS, Engberink MF, van Rooij FJ, Hofman A, Geleijnse JM. Dairy products and the risk of stroke and coronary heart disease: the Rotterdam Study. *Eur J Nutr*. 2015;54:981–90.

S19. Wang C, Yatsuya H, Tamakoshi K, Iso H, Tamakoshi A. Milk drinking and mortality: findings from the Japan collaborative cohort study. *J Epidemiol*. 2015; 25: 66–73.

S20. Dik VK, Murphy N, Siersema PD, Fedirko V, Jenab M, Kong SY, Hansen CP, Overvad K, Tjønneland A, Olsen A, Dossus L, Racine A, Bastide N, Li K, Kühn T, Boeing H, Aleksandrova K, Trichopoulou A, Trichopoulos D, Barbitsioti A, Palli D, Contiero P, Vineis P, Tumino R, Panico S, Peeters PHM, Weiderpass E, Skeie G, Hjartåker A, Amiano P, Sánchez MJ, Fonseca-Nunes A, Barricarte A, Chirlaque MD, Redondo ML, Jirström K, Manjer J, Nilsson LM, Wennberg M, Bradbury KE, Khaw KT, Wareham N, Cross AJ, Riboli E, Bueno-de-Mesquita H. Prediagnostic intake of dairy products and dietary calcium and colorectal cancer survival-results from the EPIC cohort study. *Cancer Epidemiol Biomarkers Prev*. 2014; cebp-0172.

Supplementary Data

- S21. Yang B, McCullough ML, Gapstur SM, Jacobs EJ, Bostick RM, Fedirko V, Flanders WD, Campbell PT. Calcium, vitamin D, dairy products, and mortality among colorectal cancer survivors: The Cancer Prevention Study-II Nutrition Cohort. *J Clin Oncol*. 2014; 32: 2335–2343.
- S22. Fraser GE¹, Sabaté J, Beeson WL, Strahan TM. A possible protective effect of nut consumption on risk of coronary heart disease. the adventist health study. *Arch Intern Med* 1992; 152:1416–24.
- S23. Whiteman D, Muir J, Jones L, Murphy M, Key T. Dietary questions as determinants of mortality: The OXCHECK experience. *Public Health Nutr*. 1999; 2: 477–487.
- S24. Sluik D, Boeing H, Li K, Kaaks R, Johnsen NF, Tjønneland A, Arriola L, Barricarte A, Masala G, Grioni S, et al. Lifestyle factors and mortality risk in individuals with diabetes mellitus: are the associations different from those in individuals without diabetes? *Diabetologia* 2014;57:63–72.
- S25. Boggs DA, Ban Y, Palmer JR, Rosenberg L. Higher diet quality is inversely associated with mortality in African-American women. *J Nutr* 2015;145:547–54.
- S26. Bongard V, Arveiler D, Dallongeville J, Ruidavets JB, Wagner A, Simon C, Marecaux N, Ferrieres J. Food groups associated with a reduced risk of 15-year all-cause death. *Eur J Clin Nutr* 2016;70:715–22.
- S27. Fraser GE¹, Sabaté J, Beeson WL, Strahan TM. A possible protective effect of nut consumption on risk of coronary heart disease. the adventist health study. *Arch Intern Med* 1992; 152:1416–24.
- S28. Hays JC, Keller HH, Ostbye T. The effects of nutrition-related factors on four-year mortality among a biracial sample of community-dwelling elders in the North Carolina piedmont. *J Nutr Elder* 2005; 25:41–67.

Supplementary Data

S29. Kouris-Blazos A, Gnardellis C, Wahlqvist ML, Trichopoulos D, Lukito W, Trichopoulou A. Are the advantages of the Mediterranean diet transferable to other populations? A cohort study in Melbourne, Australia. *Br J Nutr* 1999;82:57–61.

S30. Kurotani K, Akter S, Kashino I, Goto A, Mizoue T, Noda M, Sasazuki S, Sawada N, Tsugane S. Quality of diet and mortality among Japanese men and women: Japan Public Health Center based prospective study. *BMJ* 2016;352:i1209.

S31. Martínez-González MA, Guilleín-Grima F, De Irala J, Ruiz-Canela M, Bes-Rastrollo M, Beunza JJ, López del Burgo C, Toledo E, Carlos S, Sánchez-Villegas A. The Mediterranean diet is associated with a reduction in premature mortality among middle-aged adults. *J Nutr* 2012;142:1672–8.

S32. Prinelli F, Yannakoulia M, Anastasiou CA, Adorni F, Di Santo SG, Musicco M, Scarmeas N, Correa Leite ML. Mediterranean diet and other lifestyle factors in relation to 20-year all-cause mortality: a cohort study in an Italian population. *Br J Nutr* 2015;113:1003–11.

S33. Trichopoulou A, Bamia C, Trichopoulos D. Anatomy of health effects of Mediterranean diet: Greek EPIC prospective cohort study. *BMJ* 2009;338:b2337.

S34. Vormund K, Braun J, Rohrmann S, Bopp M, Ballmer P, Faeh D. Mediterranean diet and mortality in Switzerland: an alpine paradox? *Eur J Nutr* 2015;54:139–48.

S35. Yu D, Sonderman J, Buchowski MS, McLaughlin JK, Shu XO, Steinwandel M, Signorello LB, Zhang X, Hargreaves MK, Blot WJ, et al. Healthy eating and risks of total and cause-specific death among low-income populations of African-Americans and other adults in the southeastern United States: a prospective cohort study. *PLoS Med* 2015;12:e1001830.

Supplementary Data

S36. Yu D, Zhang X, Xiang YB, Yang G, Li H, Gao YT, Zheng W, Shu XO. Adherence to dietary guidelines and mortality: a report from prospective cohort studies of 134,000 Chinese adults in urban Shanghai. *Am J Clin Nutr* 2014;100:693–700.